

---

---

## Utility of High-Performance Liquid Chromatography for Identification of Mycobacterial Species Rarely Encountered in Clinical Laboratories

E. Tortoli<sup>1\*</sup>, A. Bartoloni<sup>2</sup>, C. Burrini<sup>3</sup>,  
A. Mantella<sup>2</sup>, M.T. Simonetti<sup>1</sup>

High-performance liquid chromatography (HPLC) has been demonstrated to be a suitable technique for determining the species of mycobacteria on the basis of their mycolic acid pattern. Representative HPLC profiles, which are needed for the visual recognition of chromatograms, have been published for the most frequently encountered mycobacterial species. No extensive study has been reported for less common spe-

cies, and only a few, scattered chromatographic patterns are available in literature. This study evaluates the utility of this technique for the identification of several rare species. *Mycobacterium celatum*, *Mycobacterium genavense* and *Mycobacterium simiae* chromatographic profiles have been verified, and previously unreported profiles of other species investigated. The chromatographic pattern of *Mycobacterium malmoense* is presented for the first time.

With the exception of the few species for which commercial genetic probes (Gen-Probe, USA) are available, the identification of nontuberculous mycobacteria still relies on the use of cumbersome conventional tests, which take from two to six weeks. Analysis of mycobacterial fatty acids reveals a variety of high molecular-weight  $\alpha$ -alkyl,  $\beta$ -hydroxy acids (mycolic acids), whose distribution in the cell wall is species-related; their characterization has been used for the identification to the species level of such organisms (1-4). The potential of high-performance liquid chromatography (HPLC) for elucidating the distribution of mycolic acids, after they have been converted to bromophenacyl esters, has been well documented. This system has been demonstrated as adequate for identification of the most frequently encountered mycobacteria, for whom the representative patterns have been published, by means of visual or computer-aided comparison of relevant profiles (5-11). As the method is at present well standardized, we tested its usefulness for the identification of the less frequently encountered mycobacteria, which at times constitute a challenge even for the experts.

**Materials and Methods.** Part of our laboratory collection of nontuberculous mycobacteria, seven reference strains and several recent clinical isolates (Table 1), were subjected to the HPLC procedure developed at the Centers for Disease Control (CDC) (7). The identification of all the strains was achieved using conventional methods (12) and was confirmed using nucleic acid hybridization for all species for which genetic probes are commercially available. For ambiguously identified strains, the nucleotide sequence of 16S rRNA was also determined (13). For each strain, a 10- $\mu$ l loopful of colonies recently grown on Middlebrook 7H11 medium was saponified in ethanolic potassium hydroxide (25 % KOH in 50 % ethanol) in an autoclave for 1 h at 121°C. After acidification with 18.5 % HCl, mycolic

<sup>1</sup>Bacteriological and Virological Laboratory, Careggi Hospital, viale Pieraccini 24, 50139 Firenze, Italy.

<sup>2</sup>Institute of Infectious Diseases, University of Florence, 50121 Firenze, Italy.

<sup>3</sup>Institute for Clinical Researches, 50129 Firenze, Italy.

acids were extracted with chloroform and derivatized to their UV-adsorbing p-bromophenacyl esters (14); after a further acidification a chloroform extraction was finally performed.

One hundred  $\mu$ l of the extract was spiked with 5  $\mu$ l of a high molecular-weight standard (Ribi; ImmunoChem, USA), diluted to 0.5 mg/ml and 5  $\mu$ l were injected into the reverse phase C-18 ultra-sphere-XL cartridge column of an HPLC System Gold model (Beckman, USA), equipped with a 166 model detector set at 254 nm. The elution conditions changed linearly, from a 98 % methanol, 2 % dichloromethane mixture to an 80 %, 20 % mixture in 1 min and to a 35 %, 65 % mixture over the next 9 min. (flow rate = 2.5 ml/min). At the end of the run the column was reequilibrated for 2 min to the initial conditions.

Peaks were identified on the basis of their retention times, relative to that of the internal standard (RRT) and were labelled as suggested by the CDC (11).

The bromophenacyl ester profiles were matched visually with those of all the species whose representative patterns had been published (5-9, 17). For rare mycobacteria, whose HPLC patterns are not available in the literature, the comparison with published chromatograms was used to check the uniqueness of their profiles. In the latter cases, the profiles obtained from reference strains or from genotypically identified isolates were taken as references for further comparisons; when multiple isolates were available the intraspecies variability of the pattern was assessed.

**Results and Discussion.** HPLC analysis was performed on 109 isolates representing 16 nontuberculous mycobacterial species (Table 1).

The visual match with published chromatograms, assisted on occasion by the comparison of the peaks' RRTs, allowed a correct identification of all the isolates belonging to the species *Mycobacterium asiaticum*, *Mycobacterium kansasii*, *Mycobacterium gordonae*, *Mycobacterium szulgai*, *Mycobacterium xenopi* and *Mycobacterium marinum*.

On the contrary, the bromophenacyl patterns of *Mycobacterium avium*, *Mycobacterium intracellulare*, *Mycobacterium paratuberculosis* and *Mycobacterium scrofulaceum* were very similar and did not always allow a distinction between species. Five isolates conventionally identified as belonging to the *Mycobacterium avium* complex but unable to hybridize with commercial DNA probes for either *Mycobacterium avium* or *My-*

*cobacterium intracellulare* (AccuProbe; Gen-Probe, USA) displayed a profile indistinguishable from those of other strains of *Mycobacterium avium* or *Mycobacterium intracellulare*. The 16S rRNA sequences, determined in Hannover by E.C. Böttger, attributed all five such isolates to the species *Mycobacterium intracellulare*.


Our five strains of *Mycobacterium simiae* yielded very characteristic chromatograms, with three easily recognizable clusters of peaks (Figure 1). Peaks arranged in three clusters were also presented by the newly described species *Mycobacterium genavense* (15, 16), which however could be recognized mainly by the lack of a terminal peak (Figure 1), as already reported (17). The identification of *Mycobacterium malmoense* was very easy, thanks to its uninterrupted sequence of peaks (Figure 1). Our two strains of the new species *Mycobacterium celatum* showed bromophenacyl ester profiles identical to the 'Mycobacterium xenopi-like' profiles already reported for a group of unidentified strains (9) which later became the basis for the description of the new species *Mycobacterium celatum*. A closer inspection revealed that the coincidence of patterns with *Mycobacterium xenopi* is limited to the first cluster of peaks, as those of the second one, although similar, can be differentiated on the basis of their RRTs (Figure 1). Unique profiles were also presented by rarely encountered species like: *Mycobacterium haemophilum*, *Mycobacterium shimoidei* and *Mycobacterium thermoresistibile*.

The use of bromophenacyl esters of mycolic acids for the identification of mycobacteria is presently a well standardized technique which gives consistent results. The system is neither quantitative nor able to determine the nature of the mycolic acids extracted from the mycobacterial cells, however it is suitable for rapid species identifications.

**Table 1:** Strains subjected to HPLC analysis.

Species	No. of strains
<i>Mycobacterium gordonae</i>	30
<i>Mycobacterium avium</i>	22
<i>Mycobacterium kansasii</i>	18
<i>Mycobacterium xenopi</i>	10
<i>Mycobacterium intracellulare</i>	8
<i>Mycobacterium simiae</i>	6
<i>Mycobacterium scrofulaceum</i>	5
<i>Mycobacterium malmoense</i>	3
<i>Mycobacterium genavense</i>	2
Others*	7

\* One strain each of the following species: *M. asiaticum*, *M. haemophilum*, *M. marinum*, *M. paratuberculosis*, *M. shimoidei*, *M. szulgai*, *M. thermoresistibile*.



**Figure 1:** Representative HPLC pattern of the bromophenacyl esters of mycolic acids. (IS: internal standard).

Reference chromatograms are so far available only for the species more frequently encountered in clinical laboratories and are not available for the rare or fastidious mycobacteria.

Although the limited number of samples available imposes some caution, this study shows that the adoption of this technique for the identification of the less frequent mycobacterial species is extremely promising, especially in view of the fact that no commercial probes are likely ever to be developed for their identification.

A limitation in the accuracy of identification was found for isolates of the *Mycobacterium avium* complex and of *Mycobacterium scrofulaceum*, which presented patterns not consistently different. For these species, the study of supplemental features is highly recommended for a correct identification.

The patterns of other strains considered in this work were all easily distinguishable between species and consistently similar within species. The simple visual inspection of chromatograms, aided by peak identification on the basis of their RRT,

allowed the identification of all the rare microorganisms tested.

Interestingly, HPLC seems less subject to variability than other characteristics. Such is the case with *Mycobacterium kansasii*, of which two biotypes have been recently reported, one of which fails to hybridize with commercial DNA probes (18); no difference in the two HPLC profiles has been observed. Similarly, the mycolic-acid pattern was uniform within a species, even when strains showed variable phenotypic traits (for example, niacin accumulation and photochromogenicity in *Mycobacterium simiae*).

In conclusion, we confirm the usefulness of HPLC profiles for the species identification of mycobacteria, and we endorse it for the less frequently encountered isolates. Even in the absence of a matching reference pattern, HPLC may help to avoid, on the basis of profile incompatibility, incorrect identifications that could spring from conventional procedures. The publication of further profiles will contribute to the diffusion of this rapid technique in mycobacteriological laboratories, as will the adoption of a

standard method for labelling the peaks; the CDC scheme (11), which we followed, seems adequate for this. The exploitation of automatic algorithms for the comparison of chromatograms will undoubtedly enhance the performance of this method (11).

### Acknowledgements

We thank E.C. Böttger (Institut für Medizinische Mikrobiologie, Medizinische Hochschule, Hannover, Germany) for genomic sequencing of several strains and P. Urbano (Institute of Microbiology, University of Florence, Florence, Italy) for the critical revision of the manuscript.

### References

- Minnikin DE, Goodfellow M: Lipid composition in the classification and identification of acid-fast bacteria. In: Goodfellow M, Board RG (ed): Microbiological classification and identification. Academic Press, London, 1980, p. 189–256.
- Minnikin DE, Hutchison IG, Caldicott AB, Goodfellow M: Thin-layer chromatography of methanolsates of mycolic acid-containing bacteria. *Journal of Chromatography* 1980, 188: 221–233.
- Minnikin DE, Minnikin SM, Hutchison IG, Goodfellow M, Grange JM: Mycolic acid pattern of representative strains of *Mycobacterium fortuitum*, *Mycobacterium peregrinum* and *Mycobacterium smegmatis*. *Journal of General Microbiology* 1984, 130: 363–367.
- Minnikin DE, Al-Shamaony L, Goodfellow M: Differentiation of *Mycobacterium*, *Nocardia*, and related taxa by thin-layer chromatographic analysis of whole-organism methanolsates. *Journal of General Microbiology* 1975, 88: 200–204.
- Butler WR, Kilburn JO: Identification of major slowly growing pathogenic mycobacteria and *Mycobacterium gordonae* by high-performance liquid chromatography of their mycolic acids. *Journal of Clinical Microbiology* 1988, 26: 50–53.
- Butler WR, Kilburn JO: High-performance liquid chromatography patterns of mycolic acids as criteria for identification of *Mycobacterium chelonae*, *Mycobacterium fortuitum*, and *Mycobacterium smegmatis*. *Journal of Clinical Microbiology* 1990, 28: 2094–2098.
- Butler WR, Jost KC, Kilburn JO: Identification of mycobacteria by high-performance liquid chromatography. *Journal of Clinical Microbiology* 1991, 29: 2468–2472.
- Cage GD: High-performance liquid chromatography patterns of *Mycobacterium gordonae* mycolic acids. *Journal of Clinical Microbiology* 1992, 30: 2402–2407.
- Butler WR, Thibert L, Kilburn JO: Identification of *Mycobacterium avium* complex strains and some similar species by high-performance liquid chromatography. *Journal of Clinical Microbiology* 1992, 30: 2698–2704.
- Thibert L, Lapierre S: Routine application of high-performance liquid chromatography for identification of mycobacteria. *Journal of Clinical Microbiology* 1993, 31: 1759–1763.
- Glickman SE, Kilburn JO, Butler WR, Ramos LS: Rapid identification of mycolic acid patterns of mycobacteria by high-performance liquid chromatography using pattern recognition software and a *Mycobacterium* library. *Journal of Clinical Microbiology* 1994, 32: 740–745.
- Roberts GD, Koneman EW, Kim YK: *Mycobacterium*. In: Balows A, Hausler WJ, Herrmann KL, Isenberg HD, Shadomy HJ (ed): Manual of clinical microbiology. American Society for Microbiology, Washington, DC, 1991, p. 304–339.
- Kirschner P, Springer B, Vogel U, Meier A, Wrede A, Kiekenbeck M, Bange FC, Böttger EC: Genotypic identification of mycobacteria by nucleic acid sequence determination. Report of a 2-year experience in a clinical laboratory. *Journal of Clinical Microbiology* 1993, 31: 2882–2889.
- Durst HD, Milano M, Kikta EJ, Connelly SA, Grushka E: Phenacyl esters of fatty acids via crown ether catalysts for enhanced ultraviolet detection in liquid chromatography. *Analytical Chemistry* 1975, 47: 1797–1801.
- Böttger EC, Teske A, Kirschner P, Bost S, Chang HR, Beer V, Hirschel B: Disseminated "*Mycobacterium genavense*" infection in patients with AIDS. *Lancet* 1992, 340: 76–80.
- Tortoli E, Simonetti MT, Dionisio D, Meli M: Cultural studies on two isolates of "*Mycobacterium genavense*" from patients with acquired immunodeficiency syndrome. *Diagnostic Microbiology and Infectious Disease* 1994, 18: 7–12.
- Coyle MB, Carlson LDC, Wallis CK, Leonard RB, Raisys VA, Kilburn JO, Samadpour M, Böttger EC: Laboratory aspects of "*Mycobacterium genavense*", a proposed species isolated from AIDS patients. *Journal of Clinical Microbiology* 1992, 30: 3206–3212.
- Tortoli E, Simonetti MT, Lacchini C, Penati V, Urbano P: Tentative evidence of AIDS-associated biotype of *Mycobacterium kansasii*. *Journal of Clinical Microbiology* 1994, 32: 1779–1782.