

Consensus management recommendations for less common non-tuberculous mycobacterial pulmonary diseases



Christoph Lange, Erik C Böttger, Emmanuelle Cambau, David E Griffith, Lorenzo Guglielmetti, Jakko van Ingen, Shandra L Knight, Theodore K Marras, Kenneth N Olivier, Miguel Santin, Jason E Stout, Enrico Tortoli, Dirk Wagner, Kevin Winthrop, Charles L Daley, on behalf of the expert panel group for management recommendations in non-tuberculous mycobacterial pulmonary diseases*

The 2020 clinical practice guideline for the treatment of non-tuberculous mycobacterial pulmonary disease (NTM-PD) by the American Thoracic Society, European Respiratory Society, European Society of Clinical Microbiology and Infectious Diseases, and Infectious Diseases Society of America; and the 2017 management guideline by the British Thoracic Society covered pulmonary diseases in adults caused by *Mycobacterium avium* complex, *Mycobacterium kansasii*, *Mycobacterium xenopi*, and *Mycobacterium abscessus*. In order to provide evidence-based recommendations for the treatment of less common non-tuberculous mycobacterial (NTM) species in adult patients without cystic fibrosis or HIV infection, our expert panel group performed systematic literature searches to provide management guidance for pulmonary diseases caused by seven additional organisms: *Mycobacterium chelonae*, *Mycobacterium fortuitum*, *Mycobacterium genavense*, *Mycobacterium gordonae*, *Mycobacterium malmoense*, *Mycobacterium simiae*, and *Mycobacterium szulgai*. Treatment recommendations were developed by a structured consensus process. The evidence from the scientific literature published in English for treatment recommendations for pulmonary diseases caused by other NTM species was of very low quality, with the exception of *M malmoense*, and based on the evaluation of case reports and case series. For *M malmoense*, results from two randomised controlled trials and three retrospective cohort studies provided a better evidence base for treatment recommendations, although the evidence was still of low quality.

Introduction

The 2020 updated management guideline for patients with non-tuberculous mycobacterial pulmonary diseases (NTM-PD) focuses on population, intervention, comparison, and outcome question-guided management recommendations for pulmonary disease in adults caused by *Mycobacterium avium* complex, *Mycobacterium kansasii*, *Mycobacterium xenopi*, and *Mycobacterium abscessus*.^{1,2} However, management options for NTM-PD caused by other clinically relevant non-tuberculous mycobacteria (NTM) covered in the previous management guideline are also needed for the care of affected patients.³ At present, treatment recommendations for patients with other NTM-PDs are primarily based on expert opinions and often variable. Health-care providers of patients with NTM-PDs should consult with a clinical microbiologist who has expertise in identification and antimycobacterial drug susceptibility testing, and a clinician with expertise in managing NTM disease. However, evidence-based management decisions are needed for patients affected by NTM not covered in the 2017 guideline by the British Thoracic Society (BTS) or the 2020 American Thoracic Society (ATS), European Respiratory Society (ERS), European Society of Microbiology and Clinical Infectious Diseases (ESCMID), and Infectious Diseases Society of America (IDSA) guideline.⁴

The panel of members of the 2020 ATS, ERS, ESCMID, and IDSA guideline committee did systematic reviews of the literature, independently of the societies involved in the original task force, to provide management guidance for pulmonary diseases caused by seven additional organisms: *Mycobacterium chelonae*, *Mycobacterium fortuitum*, *Mycobacterium genavense*,

Mycobacterium gordonae, *Mycobacterium malmoense*, *Mycobacterium simiae*, and *Mycobacterium szulgai*.

The following consensus guidance includes recommendations for antibiotics guided by in-vitro susceptibility results. With the exception of *M chelonae* and *M fortuitum*, there are no validated break points defining susceptibility and resistance for any of the antibiotics used for the organisms considered here.⁵

Methods

A search (see appendix pp 4–7) was adapted for execution on the Ovid MEDLINE platform and Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Embase, and the Cochrane Central Register of Controlled Trials. Searches were limited to human studies or studies indexed with neither human nor animal and those published in English. A final update was executed on Jan 6, 2020. Teams of two or three independent experts evaluated the search results for eligibility; they supplemented the electronic search by contacting experts and handsearching journals, conference proceedings, reference lists, and regulatory agency websites for relevant articles.

In addition to the systematic reviews, the members of the panel ascertained agreement on management options in a six-step consensus process as previously published.^{6,8} The results of the panel experts' votes, either in favour or against the treatment approach, is presented in the appendix (appendix p 1).

General recommendations

The expert panel recommends that health-care providers consult with a clinical microbiologist who has expertise

Lancet Infect Dis 2022

Published Online
January 25, 2022
[https://doi.org/10.1016/S1473-3099\(21\)00586-7](https://doi.org/10.1016/S1473-3099(21)00586-7)

*Members listed in the appendix

Division of Clinical Infectious Diseases, Research Center Borstel, Borstel, Germany (Prof C Lange MD); German Center for Infection Research (DZIF), Respiratory Medicine & International Health, University of Lübeck, Lübeck, Germany (C Lange); Baylor College of Medicine, Houston, TX, USA (C Lange); Institute of Medical Microbiology, National Reference Center for Mycobacteria, University of Zurich, Zurich, Switzerland (Prof E C Böttger MD); National Reference Center for Mycobacteria and Antimicrobial Resistance, AP-HP GHU Nord, Mycobacteriology, Inserm, IAME UMR 1137, University of Paris, Paris, France (Prof E Cambau MD); National Jewish Health and University of Colorado Health Sciences (Prof D E Griffith MD, Prof C L Daley MD, L Guglielmetti MD) and Library and Knowledge Services (S L Knight MS), Denver, CO, USA; Sorbonne University, Inserm, Centre for Immunology and Microbial Infections, Paris, France (L Guglielmetti); Radboud Center for Infectious Diseases, Department of Medical Microbiology, Radboud University Medical Center, Nijmegen, Netherlands (Prof J van Ingen MD); Department of Medicine, University of Toronto and University Health Network, Toronto, ON, Canada (T K Marras MD); Pulmonary Branch, National Heart, Lung and Blood Institute, Bethesda, MD, USA (K N Olivier MD); Service of Infectious Diseases, Bellvitge University Hospital-IDIBELL, University of Barcelona, L'Hospitalet de Llobregat, Barcelona, Spain

(M Santin MD); Division of Infectious Diseases and International Health, Duke University Medical Center, Durham, NC, USA

(Prof J E Stout MD); Emerging Bacterial Pathogens Unit, IRCCS San Raffaele Scientific Institute, Milan, Italy

(E Tortoli ScD); Division of Infectious Diseases, Department of Medicine II, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany (Prof D Wagner MD); Divisions of Infectious Diseases, Schools of Public Health and Medicine, Oregon Health and Science University, Portland, OR, USA

(Prof K Winthrop MD)

Correspondence to: Prof Christoph Lange, Division of Clinical Infectious Diseases, Research Center Borstel, 23845 Borstel, Germany clange@fz-borstel.de

See Online for appendix

Panel: Clinical and microbiological criteria for diagnosis of non-tuberculous mycobacteria

Clinical

Pulmonary or systemic symptoms; and nodular or cavitary opacities on chest radiograph or a high-resolution computed tomography scan that shows bronchiectasis with multiple small nodules and appropriate exclusion of other diagnoses.

Microbiological*

- Positive culture results for a non-tuberculous mycobacteria (NTM) from at least two separate expectorated sputum samples. If the results are nondiagnostic, consider repeat sputum acid-fast bacilli smears and cultures.
- or
- Positive culture results for a NTM from at least one bronchial wash or lavage.
- or
- Transbronchial or other lung biopsy with mycobacterial histologic features (granulomatous inflammation or acid-fast bacilli) and positive culture for NTM or biopsy showing mycobacterial histologic features (granulomatous inflammation or acid-fast bacilli) and one or more expectorated sputum or bronchial washings that are culture positive for NTM.

Expert consultation should be obtained when NTM are recovered that are either infrequently encountered or that usually represent environmental contamination. Patients who are suspected of having NTM pulmonary disease but do not meet the diagnostic criteria should be followed up until the diagnosis is firmly established or excluded. Making the diagnosis of NTM pulmonary disease does not per se, necessitate the institution of therapy, which is a decision based on the potential risks and benefits of therapy for individual patients.

*When two positive cultures are obtained, the isolates should be the same NTM species in order to meet disease criteria.

in identification and antimycobacterial drug susceptibility testing. Patients should be referred to a clinician with expertise in managing NTM-PD before treatment against any of the mycobacterial pathogens described in this consensus document is initiated. The clinical and microbiological criteria for the diagnosis of NTM-PD are displayed in panel.

The choice of treatment regimen should be guided by the results of the antimycobacterial drug susceptibility testing, although for some species causing NTM-PD—ie, *M genavense* and *M simiae*—the correlation of antimycobacterial drug susceptibility testing and clinical outcome can be poor (figure). For orientation, the minimum inhibitory concentrations (MIC) of antibiotics against select NTM species required to inhibit the growth of 50% and 90% of organisms (MIC₅₀ and MIC₉₀) are

provided in the appendix (appendix p 2). The major contributions of the mycobacteriology laboratory are to correctly identify the NTM species, perform antimycobacterial drug susceptibility testing, and detect acquired and inducible resistance. The dosing of drugs for the treatment of NTM-PDs mentioned in this document follows recent ATS, ERS, ESCMID, and IDSA recommendations (table 1).^{1,2} Treatment considerations are summarised in tables 2 and 3 and the quality of the evidence and strength of the recommendations are displayed in the appendix (appendix p 3)

To monitor treatment (supplement table 2, appendix p 3), sputum samples should be collected for culture every 1–2 months after initiation of therapy until there is sputum conversion to culture negative (at least two consecutive negative mycobacterial sputum cultures, collected at least 4 weeks apart during antibiotic treatment); the sampling date of the first negative culture is the date of sputum conversion to culture negative.⁶ Thereafter, sputum should be collected every 2–3 months until therapy is completed, defined as 12 months of negative mycobacterial sputum cultures (while on therapy) from the date of the first negative culture.

Rapidly growing NTM

M chelonae

M chelonae is an unusual cause of NTM-PD;^{3,9,10} therefore, clinicians should carefully assess for other causes of the patients symptoms and verify fulfillment of disease criteria before embarking on a course of treatment. Isolates are usually susceptible to tobramycin, macrolides (eg, clarithromycin and azithromycin), clofazimine, linezolid, and sometimes to fluoroquinolones and imipenem.^{3,11–15} Tobramycin is more active in vitro than amikacin and *M chelonae* is generally resistant to cefoxitin.¹⁶ *M chelonae* does not contain an erythromycin resistance methylase (*erm*) gene,¹⁷ so the macrolides are considered fully active, unlike *M fortuitum* and some *M abscessus* subspecies, which have functional *erm* genes. Antimycobacterial drug susceptibility testing should be done, as well as detection of acquired resistance to macrolides, which was described in this species, although mostly in non-pulmonary infections.¹⁸

A systematic literature review by two independent experts identified 18 case reports and case series describing 57 patients with *M chelonae* pulmonary disease in the scientific literature published in English, but no evidence-based management recommendations for *M chelonae* pulmonary disease based on clinical trial data.^{9,14,19–34} In most patients, risk factors for *M chelonae* pulmonary disease were not stated or could not be identified. Approximately 15% of patients who have been reported with *M chelonae* pulmonary disease were post-organ transplant.^{19,21,23} By contrast with *M fortuitum* pulmonary disease, gastrointestinal motility disorders (eg, achalasia) did not appear to be a prominent risk

factor for *M chelonae* pulmonary disease. The clinical spectrum of *M chelonae* pulmonary disease includes bronchiolitis, fibronodular or patchy consolidations, cavitary disease, and rarely empyema.

Treatment for *M chelonae* pulmonary disease initially includes at least two drugs for mild to moderate disease or three drugs in more severe disease that the organism shows in-vitro susceptibility to. Treatment often includes one or two intravenous drugs to be continued for an initial period, usually for 4–16 weeks, to achieve control of disease; tobramycin is the preferred aminoglycoside. At least two oral drugs, one of which should be a macrolide that has demonstrated in-vitro susceptibility, are administered for a total duration of therapy of at least 12 months after conversion to culture negative, to complete the treatment regimen.

Favoured antibiotics include azithromycin (250–500 mg) once a day or clarithromycin (500 mg) twice a day, tobramycin (4–5–7 mg/kg) intravenously once a day or tobramycin (5–7 mg/kg) intravenously three times a week, imipenem-cilastatin (1 g each) intravenously two or three times a day or a fluoroquinolone—eg, moxifloxacin (400 mg) once a day—clofazimine (100–200 mg) once a day, or linezolid (600 mg) once a day, based on in-vitro antimycobacterial drug susceptibility test results. Caution must be exercised in the prolonged use of aminoglycosides because of the risk of otovestibular toxicity and nephrotoxicity. Approximately 20% of patients with *M chelonae* pulmonary disease received adjunctive partial pulmonary surgical resection,^{20,25} these patients generally had a successful treatment outcome, thus surgical resection should be considered when it is an option. Based on the available literature, it was not possible to state an association between the type of drug therapy against *M chelonae* pulmonary disease and successful treatment outcome.

M fortuitum

Although *M fortuitum* is a relatively common NTM species isolated from respiratory specimens, it is a relatively uncommon cause of NTM-PD.^{3,9,35} Clinicians should maintain a high-diagnostic threshold for identifying *M fortuitum* pulmonary disease, and they should carefully assess for other causes of the patients symptoms and radiological abnormalities. *M fortuitum* isolates are usually susceptible to multiple antibiotics including fluoroquinolones, doxycycline, minocycline, and sulfonamides.³ In-vitro data suggest susceptibility to clofazimine (appendix p 2). In addition, isolates are typically susceptible to amikacin and imipenem. *M fortuitum* usually contains an *erm* (39) gene which confers inducible resistance; therefore, macrolides should not be considered active.³⁶ Antimycobacterial drug susceptibility testing should be done, as well as detection of acquired resistance to fluoroquinolones, which was described in this species, although mostly in non-pulmonary infections.³⁷


A systematic literature review by two independent experts identified 45 case reports and case series describing 150 patients with *M fortuitum* pulmonary disease in the scientific literature published in English, but no evidence-based management recommendations based on clinical trial data for *M fortuitum* pulmonary disease.^{11,19,21,24,26,28,29,31,32,34,35,38–71}

M fortuitum pulmonary disease occurs predominantly in patients with underlying conditions: oesophageal and gastrointestinal motility problems (eg, achalasia),^{40,41,45} structural lung changes such as post-tuberculosis changes^{35,38,54} severe chronic obstructive pulmonary disease (COPD),^{58,69} and cancer.^{24,58} Conversion to sputum culture negative for *M fortuitum* without antibiotic treatment is not uncommon when the underlying cause (eg, achalasia) has been addressed.⁹ The underlying cause must be addressed if possible; with its correction or improvement, antibiotic treatment might not be required, or a less intensive regimen might suffice. However, *M fortuitum* pulmonary disease is associated with a poor prognosis and about a-third of the affected patients reported in the literature do not survive despite antibiotic therapy, probably reflecting the severity of the underlying disease.^{28,45,58,68,71}

Patients who received fluoroquinolones as part of their treatment regimen seemed to have a favourable treatment outcome, but we did not find any publications that met the criteria of a clinical trial for evidence-based management recommendations for *M fortuitum* pulmonary disease.^{38,42,53,54,56,58,63,67}

Clinicians should be aware that there are other mycobacterial species related to *M fortuitum* (*Mycobacterium peregrinum*, *Mycobacterium septicum*, *Mycobacterium porcinum*, and *Mycobacterium conceptionense*) included in the so-called *M fortuitum* complex, which might present different antibiotic susceptibilities, including macrolide susceptibility.⁷²

Treatment for *M fortuitum* pulmonary disease should include at least two drugs for mild-to-moderate disease or three drugs in more severe disease that the organism shows in-vitro susceptibility to. Treatment often includes one or two intravenous drugs to be continued for an initial period, usually for 4–16 weeks, to achieve control of disease; amikacin is the preferred aminoglycoside. Frequently used doses include amikacin (15–20 mg/kg) intravenously once a day or amikacin (15–25 mg/kg) three times a week in severe disease manifestations (lower in older patients and in prolonged treatment). Caution must be exercised in the prolonged use of aminoglycosides because of the risk of otovestibular toxicity and nephrotoxicity. Other favoured intravenous drugs are imipenem-cilastatin (1 g each) intravenously two or three times a day or ceftazidime (3–4 g) intravenously twice a day. At least two oral drugs that have demonstrated in-vitro susceptibility should be continued until at least 12 months after conversion to culture negative to complete the treatment regimen. Favoured oral



antibiotics include fluoroquinolones—eg, moxifloxacin (400 mg) once a day or levofloxacin (750–1000 mg) once a day—co-trimoxazole (trimethoprim 800 mg and 160 mg sulfamethoxazole) twice a day, linezolid (600 mg) once a day, clofazimine (100–200 mg) once a day, or doxycycline (100–200 mg) once a day. A summary of rapidly growing NTM treatment recommendations is shown in table 2.

Slowly growing NTM

M genavense

M genavense is a rare cause of NTM-PD.^{73–75} The organism is fastidious and does not grow on solid media, although limited growth can be observed in supplemented broth cultures; molecular detection is most reliable, thus antimycobacterial drug susceptibility testing in *M genavense* is challenging.⁷⁶ A systematic literature review by two independent experts identified five case reports and case series describing six patients with *M genavense* pulmonary disease in the scientific literature published in English, but no clinical trial data to inform evidence-based management recommendations for *M genavense* pulmonary disease.^{73,77–79} *M genavense* was first described in the setting of severe immunosuppression with disseminated disease.^{3,74,75,80}

Available data suggests that most isolates are susceptible in vitro to fluoroquinolones, clofazimine, and amikacin, although susceptibility testing has not been standardised for *M genavense* (appendix p 2).^{80,81} Case series have described treatment success with macrolide containing multi-drug regimens that typically include ethambutol plus one or two other drugs such as a rifamycin, amikacin, or a fluoroquinolone.^{74,75,82–84}

Treatment for *M genavense* pulmonary disease includes at least three drugs that the organism shows in-vitro susceptibility to. The panel suggests the following treatment regimen: azithromycin (250–500 mg) once a day, rifampicin (600 mg) once a day, and ethambutol (15 mg/kg) once a day. In case of intolerance or drug resistance to macrolides, rifamycins, or ethambutol, the following treatment regime can be used instead:

moxifloxacin (400 mg) once a day, amikacin (15–20 mg/kg) intravenously once a day or amikacin (15–25 mg/kg) three times a week (lower in older patients and prolonged use), or clofazimine (100–200 mg) once a day. Treatment should be continued until at least 12 months post-conversion to culture negative to complete the treatment regimen.

M gordonae

M gordonae is known as a non-pathogenic species but is one of the most common NTM isolated from respiratory samples; probably because it is prevalent in the environment, especially the water supply network.^{85–89} It is also easily isolated in mycobacteriology laboratories (naturally orange colonies growing on all media after 2 weeks incubation). Limited data suggest that *M gordonae* is variably susceptible in vitro to several agents: clarithromycin, ciprofloxacin, linezolid, and amikacin.^{90–92} The correlation between these in-vitro results and patient outcomes is unknown.

A systematic literature review by two independent experts identified nine case reports and case series describing 13 non-immunosuppressed adults with *M gordonae* pulmonary disease in whom treatment and outcomes were adequately described, but no evidence-based management recommendations for *M gordonae* pulmonary disease based on clinical trial data.^{69,93–100}

Treatment for *M gordonae* pulmonary disease before 1990 often included isoniazid, rifampicin, and ethambutol. Treatment in newer studies more frequently included clarithromycin, rifampicin, and ethambutol. Information regarding cure was provided for 12 cases and was definite or likely in 11 (92%) patients overall and 11 patients who did not experience respiratory failure.^{69,93–100} Surgical resection, in addition to antibiotic therapy, was described in two patients, both of whom were cured.^{97,100}

When *M gordonae* is cultured in respiratory samples, other causes of lung disease should be carefully considered. Multiple sputum samples should be collected over several weeks or months to identify the presence of any other NTM known to be more pathogenic. Clinicians should maintain a high-diagnostic threshold for identifying *M gordonae* pulmonary disease. In addition to the clinical and radiological criteria for *M gordonae* pulmonary disease (as *M gordonae* is almost always non-pathogenic) described in the international NTM guidelines,⁹⁷ more than two positive sputum cultures and at least one detection of acid-fast bacilli in a respiratory specimen with documented *M gordonae* cultures should be present.

Even when current diagnostic criteria are met, treatment is seldom necessary. In the rare situation when treatment is necessary, antimycobacterial drug susceptibility testing can be done as for other NTM; although no standard treatment can be recommended, a combination of a macrolide, rifampicin, and ethambutol has been described as successful in some cases.

Figure: Optimal treatment pathways for less commonly encountered NTM-PD

The flow chart provides information on the recommended treatment for seven less common NTM. The seven species of NTM are displayed in the left-hand column of the flow chart; species coloured in green indicate high pathogenicity and high probability of causing disease, yellow indicates species which are less pathogenic with a lower probability of causing disease, and red indicated a species that likely represents a contaminant. Drugs used to treat the NTM are displayed to the right of the flow chart; green indicates drugs that are most strongly recommended, yellow indicates less certainty due to missing clinical data, orange indicates drugs that have to be given intravenously, thus are not preferred, and red indicates drugs that should not be used. NTM= non-tuberculous mycobacteria. NTM-PD=non-tuberculous mycobacterial pulmonary disease. ATS=American Thoracic Society. ERS=European Respiratory Society. ESCMID=European Society of Clinical Microbiology and Infectious Diseases. IDSA=Infectious Disease Society of America. Please see appendix for additional information (appendix p 8).

	Daily dose	Three times a week dose	Hepatic impairment dose	Renal impairment dose
Oral				
Azithromycin	250–500 mg once a day	500 mg once a day	NA	NA
Ciprofloxacin	500–750 mg twice a day	NA	NA	250–500 mg dosed at intervals according to CrCl
Clarithromycin	500 mg twice a day	500 mg once a day	NA	Reduce dose by 50% if CrCl less than 30 mL/min
Clofazimine*	100–200 mg once a day	NA	Caution in severe hepatic impairment	NA
Doxycycline	100 mg once a day or twice a day	NA	NA	NA
Ethambutol	15 mg/kg once a day	25 mg/kg once a day	NA	Increase dosing interval (eg, 15–25 mg/kg, three times a week)
Levofloxacin	750–1000 mg once a day	NA	NA	Increase dosing interval (eg, 10–15 mg/kg, three times a week)
Linezolid	600 mg once a day or twice a day†	NA	NA	NA
Moxifloxacin	400 mg once a day	NA	NA	NA
Rifabutin	150–300 mg once a day (150 mg once a day with clarithromycin)	300 mg once a day	Caution	Reduce dose by 50% if CrCl less than 30 mL/min
Rifampicin	10 mg/kg (450 mg or 600 mg) once a day	600 mg once a day	Caution	NA
Co-trimoxazole	800 mg trimethoprim and 160 mg sulfamethoxazole twice a day	NA	Caution	Reduce dose by 50% if CrCl 15–30 mL/min
Parenteral				
Amikacin‡ (IV)	15–20 mg/kg once a day§, adjusted according to drug level monitoring¶	15–25 mg/kg once a day§, adjusted according to drug level monitoring¶	NA	Reduce dose or increase dosing interval (eg, 15 mg/kg, twice a week or three times a week)
Cefoxitin (IV)	2–4 g twice a day or three times a day (maximum daily dose is 12 g/day)	NA	NA	Reduce dose or increase dosing interval
Imipenem-cilastatin (IV)	1g each, twice a day or three times a day	NA	NA	Reduce dose or increase dosing interval
Tobramycin‡ (IV)	4.5–7 mg/kg once a day	5–7 mg/kg once a day	NA	Reduce dose or increase dosing interval (eg, 5–7 mg/kg, twice a week or three times a week); inhalative route of administration not studied
<p>NA=not applicable. NTM-PD= non-tuberculous mycobacterial pulmonary disease. CrCl= creatinine clearance. IV=intravenously. This table should be considered as a general guide for dosing in clinical practice, and it does not replace specific local, national, or regional dosing guidelines. *Clofazimine availability varies by country. In the USA, an investigational new drug application is required. †Most experts recommend once a day dosing of linezolid because of the high rate of drug-related adverse reactions associated with twice a day dosing. ‡Caution must be exercised in the prolonged use of aminoglycosides because of the risk of otovestibular toxicity and nephrotoxicity. §The use of the described regimens for 15 weeks was associated with permanent ototoxicity in approximately a-third of patients, and the risk was associated with age and cumulative dose. Given the high rates of ototoxicity, risks and benefits should be carefully considered with the aims of therapy. Clinicians should consider lower dose ranges and probably rely on intermittent dosing when more prolonged therapy is employed. ¶Drug level monitoring: trough less than 5 mg/L, peak with daily dosing 35–45 µg/mL, peak with intermittent dosing 65–80 µg/mL.</p>				
Table 1: Dosing guidelines for drugs used in the management of NTM-PD				

M malmoense

M malmoense was first described in four patients from Malmö and Lund, Sweden,¹⁰¹ in 1977 and is one of the most common species causing NTM-PD in regions of northern Europe. Multiple studies using different laboratory methods have described the susceptibility of *M malmoense* to rifampicin, rifabutin, clarithromycin, and clofazimine, with variable resistance to amikacin, ethambutol, and fluoroquinolones, and resistance to isoniazid.^{102–106}

M malmoense most commonly causes pulmonary disease but can also cause cervical lymphadenitis,

tenosynovitis, skin infections, and rarely disseminated disease in the severely immunocompromised host. It is seldom isolated from the environment but it can be found in natural water and soil in some geographical areas.^{107–109} The clinical presentation of *M malmoense* pulmonary disease frequently mimics tuberculosis, with cavities and airspace disease being the most common radiological findings.^{110,111} The majority of affected patients are male, approximately half have underlying COPD or a history of pulmonary tuberculosis. When *M malmoense* is isolated from respiratory samples, it is usually of clinical significance, although there is some geographical

	Number of drugs	Drugs	Duration of therapy	Comments
<i>M chelonae</i>	Initial phase (≥3); continuation phase (≥2)	Azithromycin (once a day) or clarithromycin (twice a day); tobramycin IV (once a day or three times a week); imipenem-cilastatin (two to three times a day); moxifloxacin (once a day) or levofloxacin (once a day), or ciprofloxacin (twice a day); linezolid (once a day); clofazimine (once a day)	12 months beyond culture conversion	Very low level of evidence; drugs should be selected according to DST results when available; caution patients about aminoglycoside ototoxicity and nephrotoxicity; for mild to moderate disease an oral two-drug regimen could suffice, provided that DST has proven two such drugs to be active
<i>M fortuitum</i>	Initial phase (≥3); continuation phase (≥2)	Moxifloxacin (once a day) or levofloxacin (once a day), or ciprofloxacin (twice a day); amikacin IV (once a day or three times a week); imipenem-cilastatin (two to three times a day); cefoxitin (two to three times a day); linezolid (once a day); co-trimoxazole (twice a day); (clofazimine [once a day]); (doxycycline [twice a day])*	12 months beyond culture conversion	Very low level of evidence; drugs should be selected according to DST results when available; the detection and management of underlying oesophageal disorders or aspiration is critical; fluoroquinolones are probably the most effective; there is natural resistance to macrolides; caution patients about ototoxicity and nephrotoxicity of aminoglycosides; for mild to moderate disease an oral two-drug regimen could suffice provided that DST has proven two such drugs to be active

NTM-PD=non-tuberculous mycobacterial pulmonary disease. IV=intravenously. DST=drug susceptibility testing.*Drugs in brackets are non-preferred options.

Table 2: Treatment regimens for rapidly growing NTM-PD

	Number of drugs	Drugs*	Duration of therapy	Comments
<i>M genavense</i>	≥3	Azithromycin (once a day) or clarithromycin [twice a day]; rifampicin (once a day); ethambutol (once a day); (moxifloxacin [once a day]); (clofazimine [once a day]); (amikacin IV [once a day])	12 months beyond culture conversion	Very low level of evidence; drugs should be selected according to DST results when available; moxifloxacin or amikacin may be used in cases of intolerance or drug resistance to macrolides, rifamycins, or ethambutol
<i>M malmoense</i>	≥3	Azithromycin (once a day) or clarithromycin [twice a day]; rifampicin (once a day); ethambutol (once a day); (amikacin IV [once a day or three times a week]); (moxifloxacin [once a day]); (clofazimine [once a day])	12 months beyond culture conversion	Low level of evidence; drugs should be selected according to DST results, when available; moxifloxacin or clofazimine can be used in case of intolerance or drug resistance to macrolides, rifamycins, or ethambutol; amikacin should be added for cavitory or severe disease; caution patients about aminoglycoside ototoxicity and nephrotoxicity
<i>M szulgai</i>	≥3	Azithromycin (once a day) or clarithromycin [twice a day]; rifampicin (once a day); ethambutol (once a day); (moxifloxacin [once a day]); (clofazimine [once a day]); (amikacin IV [once a day or three times a week])	12 months or 12 months beyond culture conversion if treatment with a macrolide, a rifamycin, or ethambutol cannot be used.	Very low level of evidence; drugs should be selected according to DST results, when available; fluoroquinolones (moxifloxacin or levofloxacin), clofazimine, or aminoglycosides (streptomycin or amikacin) can be used in case of intolerance or drug resistance to macrolides, rifamycins, or ethambutol; caution patients about ototoxicity and nephrotoxicity of aminoglycosides
<i>M simiae</i>	≥3	Azithromycin or clarithromycin; moxifloxacin (once a day); co-trimoxazole (twice a day); clofazimine (once a day); (amikacin IV [once a day or three times a week])	12 months beyond culture conversion	Very low level of evidence; drugs should be selected according to DST results, when available; amikacin should be added for cavitory or severe disease; possible combinations include azithromycin, moxifloxacin, and co-trimoxazole; azithromycin, clofazimine, and amikacin; or azithromycin and moxifloxacin in combination with one or two additional drugs based on DST results with clofazimine and amikacin being the most suitable options
<i>M gordonae</i> †	NA	NA	NA	NA

IV=intravenously. DST=drug susceptibility testing. NA=not applicable. *Drugs in brackets are non-preferred options. †In general, isolation of *M gordonae* from respiratory specimens represents transient colonisation or laboratory contamination; treatment should not be offered without overwhelming evidence that this organism is causing invasive pulmonary disease. If treatment is necessary, DST can be performed; although no standard treatment can be recommended, a combination of a macrolide, rifampicin, and ethambutol has been described as successful.

Table 3: Treatment regimens for slowly growing NTM-PD

variability suggesting that pathogenicity might vary by region.^{102,112}

A systematic literature review by two independent experts found five publications (two randomised controlled trials^{104,105} and three retrospective cohort studies^{102,113,114}) that met criteria for evidence-based management recommendations for *M malmoense* pulmonary disease. In addition, two systematic reviews were identified that addressed treatment outcomes or treatment recommendations for *M malmoense* pulmonary disease.^{4,48}

An early retrospective review¹¹³ from Cardiff, UK, described the clinical outcomes of 37 patients with *M malmoense* pulmonary disease who were treated with a

variety of non-macrolide-based treatment regimens. Those who received three drugs (isoniazid, rifampicin, and ethambutol) for 18–24 months tended to do better than those who received two drugs, or those who were treated for less than 18 months.¹¹³ Among patients treated with a BTS recommended regimen (rifampicin plus different combinations of ethambutol, isoniazid, clarithromycin, or ciprofloxacin) treatment success was achieved in 75% of patients. In a series of 14 consecutive patients with fibrocavitary *M malmoense* pulmonary disease in Edinburgh, UK, investigators reported 100% conversion to culture negative and symptom reduction after 24 months of combination rifampicin

(450–600 mg) once a day, ethambutol (15 mg/kg) once a day, and clarithromycin (500 mg) twice a day.¹¹⁴ In this study initial *M malmoense* isolates from all 14 patients were judged to be susceptible to rifampicin and resistant to isoniazid, with 23% resistant to ethambutol, 8% resistant to clarithromycin, and 46% resistant to ciprofloxacin. A retrospective review of 30 patients with *M malmoense* pulmonary disease from the Netherlands found that multi-drug treatment regimens varied considerably, although most received a macrolide in addition to ethambutol and rifampicin.¹⁰² The mean duration of therapy was 12 months, which is shorter than the duration in the BTS trials. 21 (70%) patients had a good clinical response (defined as symptomatic improvement and conversion to culture negative) to treatment, five (17%) patients had failure or relapse, and four (13%) patients died.

The BTS performed a randomised controlled trial involving 106 patients with *M malmoense* pulmonary disease; patients were treated for 2 years with either rifampicin and ethambutol or rifampicin, ethambutol, and isoniazid.^{104,105} Of 106 patients, 10% of patients had a poor clinical outcome (death due to NTM, treatment failure, or relapse); there was no significant difference between groups. Only 20 (38%) of 52 patients who received rifampicin and ethambutol, and 24 (44%) of 54 of patients who additionally received isoniazid were alive and free of relapse, and therefore considered cured after 5 years. A second BTS randomised controlled trial allocated 167 patients to 2 years of treatment with rifampicin, ethambutol, and clarithromycin; or rifampicin, ethambutol, and ciprofloxacin.¹¹⁵ Although there was no significant difference in the poor outcome rate between groups (overall, 7%), more patients receiving the rifampicin, ethambutol, and clarithromycin regimen completed treatment and were alive and cured at 5 years (38% vs 20%); however, there were more side effects in this group. In addition, the number of patients alive and cured at 5 years in the clarithromycin group was the same as that seen with rifampicin and ethambutol in the previous trial. Compared with patients with *M avium* complex and *M xenopi* pulmonary disease, who were also included in the trial, patients with *M malmoense* were significantly less likely to have a poor outcome.

Two systematic reviews were identified.^{4,48} One systematic review reported that three studies^{105,114,115} investigated treatment outcomes in patients with *M malmoense* pulmonary disease, comprising a total of 287 patients.⁴⁸ The weighted average proportion of treatment success (the proportion of patients in whom sputum conversion to culture negative was achieved at the end of therapy minus the proportion of relapses reported at the end of follow-up) was 54.4% (95% CI 34.7–73.4); however, the authors noted that the comparability between the studies and the different treatment regimens was impeded by the high all-cause mortality of 34–49.1%.

The BTS NTM guideline reviewed two randomised controlled trials^{104,115} and four case series.^{102,103,113,114} For non-cavitary disease the recommended treatment was with a macrolide, rifampicin, and ethambutol. The addition of parenteral or inhaled amikacin was recommended in cavitary disease, although there was no evidence presented that supported the use of amikacin.⁴ A macrolide-containing three-drug regimen offers the best option for treatment success based on these studies, although treatment outcomes remain suboptimal and the evidence base for treatment recommendations is low.

Treatment for *M malmoense* pulmonary disease generally includes at least three drugs: azithromycin (250–500 mg) once a day or clarithromycin (500 mg) twice a day, rifampicin (600 mg) once a day, and ethambutol (15 mg/kg) once a day. Additional drugs may include moxifloxacin (400 mg) once a day or clofazimine (100–200 mg) once a day. Based on the utility of amikacin for most NTM species (except *M chelonae*), favourable in-vitro drug susceptibility profiles, and expert opinion, the addition of parenteral amikacin can be considered in severe cases, such as cavitary disease. The optimal duration of therapy is unknown, but consistent with other NTM pulmonary pathogens the expert panel recommends a total duration of therapy of at least 12 months post-conversion to culture negative to complete the treatment regimen.

M simiae

M simiae is a rare human pathogen causing disseminated or pulmonary disease; only 4–21% of patients with respiratory *M simiae* isolates fulfil the ATS diagnostic criteria.^{116–118} The rates of isolation vary geographically¹¹⁹ with higher rates being reported in Israel^{120,121}, Lebanon¹²², Iran¹²³ and India.¹²⁴ Respiratory isolates of *M simiae* are most often contaminants, thus clinicians should maintain a high threshold to diagnose *M simiae* pulmonary disease and carefully assess for other causes of a patient's symptoms or radiological abnormalities. *M simiae* is considered among the hardest to treat NTM owing to its natural drug resistance; most antimycobacterial drug susceptibility testing reports of *M simiae* describe resistance to most tested drugs.^{116,118,125} In vitro, the *M simiae* strain ATCC 25275 was resistant to most of the 19 drugs tested, except macrolides, clofazimine, and sulfamethoxazole (quinolones and aminoglycosides were not tested).¹²⁶ Furthermore, there is in-vitro synergy between clofazimine and amikacin for *M simiae* isolates.¹¹ In addition, variable proportions of clinical strains can be susceptible to moxifloxacin (and to ciprofloxacin, to a lesser extent), aminoglycosides, and cycloserine.^{125,127}

A systematic literature review by two independent experts identified 11 case reports and case series describing 197 patients with *M simiae* pulmonary disease in the scientific literature published in English, but no evidence-based management recommendations for *M simiae* pulmonary disease based on clinical trial

data.^{116,117,121,122,128-134} The largest study of 102 patients was from Israel; no treatment failures or relapses were reported during a mean follow-up of 24 months, with patients receiving clarithromycin, ethambutol, and rifampicin treatment for at least 12 months.¹²¹ Although the 1997 ATS criteria were reportedly used to select patients, 38 (37%) of 102 patients were asymptomatic and had an unremarkable chest radiograph; therefore, not fulfilling these criteria. The expert panel concluded that data from this study could not be considered reliable enough to form a basis for treatment recommendations.¹²¹

The treatment regimen for *M simiae* pulmonary disease includes at least three drugs: azithromycin (250–500 mg) once a day or clarithromycin (500 mg) twice a day, moxifloxacin (400 mg) once a day, co-trimoxazole (trimethoprim 800 mg and sulfamethoxazole 160 mg) twice a day, amikacin (15–20 mg/kg) intravenously once a day or amikacin (15–25 mg/kg) three times a week (lower in older patients and with prolonged use), or clofazimine (100–200 mg) once a day. Depending on antimycobacterial drug susceptibility testing, accepted treatment regimens could be a clofazimine, amikacin, and azithromycin combination or a azithromycin, moxifloxacin, and co-trimoxazole combination,^{3,16} or a azithromycin and moxifloxacin combination with one or two additional drugs based on antimycobacterial drug susceptibility testing results; clofazimine and aminoglycosides would be the most appropriate options.¹²⁵ Surgical resection of affected lobes should be evaluated as an adjunctive treatment option.¹⁶ Optimal treatment duration is unknown; experts consider treating for at least 12 months after conversion to negative sputum culture, possibly introducing a step-down approach to consolidation therapy (eg, stop intravenous amikacin) once clinical, microbiological, and radiological improvement has been documented.

M szulgai

M szulgai is an unusual cause of pulmonary disease and accounts for less than 1% of human respiratory isolates of NTM; however, the isolation of *M szulgai* appears to be clinically relevant in most cases.¹³⁵⁻¹³⁷ Isolates are usually susceptible in vitro to clarithromycin and rifampicin. The largest case series reporting on antimycobacterial susceptibility testing of *M szulgai* included 23 strains of *M szulgai*.¹⁰⁶ Drug resistance was observed in all 23 (100%) strains to isoniazid, in 9% of strains to ethambutol, in 26% of strains to ciprofloxacin, 13% of strains to amikacin, and in 4% of strains to rifampicin. All 23 strains of *M szulgai* were susceptible to clarithromycin. In-vitro data suggest susceptibility to clofazimine (appendix p 2). Later generation fluoroquinolones were not tested.

A systematic literature review conducted by two independent experts identified 25 retrospective case reports and case series, including a total of 44 patients with *M szulgai* pulmonary disease described in the scientific

literature published in English^{136,138-149} but no evidence-based management recommendations for *M szulgai* pulmonary disease based on clinical trial data. Most patients were treated with a combination of rifampicin, ethambutol, and clarithromycin or azithromycin. Other treatments included a mixture of rifampicin-based regimens; usually a combination of rifampicin with two or more drugs: isoniazid, ethambutol, fluoroquinolones, ethionamide, and intravenous drugs. Treatment duration was variable; 12 months was most frequently used (range 5–18 months). The outcome was favourable in 85% of patients treated with rifampicin, clarithromycin, and azithromycin combination regimens; no relapses were observed among the five (11%) patients that post-treatment follow-up was available for. The cure rate was 81% among 21 patients treated with clarithromycin and azithromycin sparing regimens; unfavourable outcomes included one failure of treatment after 6 months and one relapse related to non-compliance. Four (9%) patients underwent surgery, but it was only performed to treat *M szulgai* pulmonary disease in two of these patients, and one of these two patients had a tissue culture sample that was negative for *M szulgai*.

Treatment for *M szulgai* pulmonary disease should include at least three drugs that the organism shows in-vitro susceptibility for. Based on the low level evidence available, a combination of rifampicin (600 mg) once a day, azithromycin (250–500 mg) once a day or clarithromycin (500 mg) twice a day, and ethambutol (15 mg/kg) once a day for 12 months is recommended. Clofazimine (100–200 mg) once a day can be used if there is intolerance or drug resistance to macrolides, rifamycins, or ethambutol. Intravenously administered drugs (eg, amikacin) are alternatives in case of intolerance or resistance to the recommended orally available drugs. There are no in-vitro susceptibility data or clinical experience with later-generation fluoroquinolones. In such cases, if a treatment with a macrolide, rifamycin, and ethambutol cannot be used, it is advised to prolong treatment to 12 months after conversion to culture negative. There is no evidence to recommend surgery as part of the treatment for *M szulgai* pulmonary disease.

A summary of slowly growing NTM treatment recommendations is shown in table 3.

Conclusion

We did a systematic review of the literature and consensus process to provide treatment recommendations for pulmonary diseases caused by seven additional NTM not covered in the recent ATS, ERS, ESCMID, and IDSA or BTS clinical practice guidelines.^{1,2,4}

With the exception of *M malmoense*, where recommendations are based on two randomised controlled trials and three retrospective cohort studies, the consensus recommendations by the panel members for the other six NTM species are based on case reports and case series only, thus are graded to be very low level evidence. Higher level evidence from patient registries

and clinical trials to determine the best treatments for patients affected by pulmonary diseases caused by the other six NTM species are needed to inform clinicians in the future about the best management options for their patients.

Contributors

CL, CLD, and EC developed the idea and outline for the manuscript. SLK performed the electronic systematic literature searches. All co-authors contributed equally to the hand-search of the selected literature, the consensus process, drafting of the manuscript, and manuscript revision.

Declaration of interests

CLD reports grants and personal fees from Insmmed, Paratek, and Spero; personal fees from AN2 and Matinas, and grants from BugWorks; outside the submitted work. DEB reports grants and personal fees from Insmmed. CL is supported by the German Center for Infection Research (DZIF) and reports personal fees from Chiesi, Gilead, Janssen, Novartis, Oxfordimmunotec, and Insmmed outside the submitted work. TKM reports grants and personal fees from Insmmed; and personal fees from Astra Zeneca, RedHill Biopharma, Novartis, and Spero; outside the submitted work. KNO is supported by the intramural research program of the NHLBI, NIH and reports grants from Beyond Air outside the submitted work. KW reports grants and personal fees from Insmmed; and personal fees from Paratek, Red Hill Biopharma, Horizon, and Spero; outside the submitted work. All other authors declare no competing interests.

Acknowledgments

CL is supported by the German Center for Infection Research (DZIF). KNO is supported by the intramural research program of The National Heart, Lung, and Blood Institute (NHLBI), National Institutes of Health (NIH).

References

- Daley CL, Iaccarino JM, Lange C, et al. Treatment of nontuberculous mycobacterial pulmonary disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Eur Respir J* 2020; **56**: 2000535.
- Daley CL, Iaccarino JM, Lange C, et al. Treatment of nontuberculous mycobacterial pulmonary disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Clin Infect Dis* 2020; **71**: 905–13.
- Griffith DE, Aksamit T, Brown-Elliott BA, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007; **175**: 367–416.
- Haworth CS, Banks J, Capstick T, et al. British Thoracic Society guidelines for the management of non-tuberculous mycobacterial pulmonary disease (NTM-PD). *Thorax* 2017; **72** (suppl 2): ii1–64.
- Brown-Elliott BA, Woods GL. Antimycobacterial susceptibility testing of nontuberculous mycobacteria. *J Clin Microbiol* 2019; **57**: e00834-19.
- van Ingen J, Aksamit T, Andrejak C, et al. Treatment outcome definitions in nontuberculous mycobacterial pulmonary disease: an NTM-NET consensus statement. *Eur Respir J* 2018; **51**: 1800170.
- Lange C, Abubakar I, Alffenaar JW, et al. Management of patients with multidrug-resistant/extensively drug-resistant tuberculosis in Europe: a TBNET consensus statement. *Eur Respir J* 2014; **44**: 23–63.
- Domínguez J, Boettger EC, Cirillo D, et al. Clinical implications of molecular drug resistance testing for *Mycobacterium tuberculosis*: a TBNET/RESIST-TB consensus statement. *Int J Tuberc Lung Dis* 2016; **20**: 24–42.
- Griffith DE, Girard WM, Wallace RJ Jr. Clinical features of pulmonary disease caused by rapidly growing mycobacteria. An analysis of 154 patients. *Am Rev Respir Dis* 1993; **147**: 1271–78.
- van Ingen J, de Zwaan R, Dekhuijzen RP, Boeree MJ, van Soolingen D. Clinical relevance of *Mycobacterium chelonae-abscessus* group isolation in 95 patients. *J Infect* 2009; **59**: 324–31.
- van Ingen J, Totten SE, Helstrom NK, Heifets LB, Boeree MJ, Daley CL. In vitro synergy between clofazimine and amikacin in treatment of nontuberculous mycobacterial disease. *Antimicrob Agents Chemother* 2012; **56**: 6324–27.
- Dalovisio JR, Pankey GA, Wallace RJ, Jones DB. Clinical usefulness of amikacin and doxycycline in the treatment of infection due to *Mycobacterium fortuitum* and *Mycobacterium chelonae*. *Rev Infect Dis* 1981; **3**: 1068–74.
- Yew WW, Kwan SY, Wong PC, Lee J. Ofloxacin and imipenem in the treatment of *Mycobacterium fortuitum* and *Mycobacterium chelonae* lung infections. *Tubercle* 1990; **71**: 131–33.
- van Ingen J, Ferro BE, Hoefsloot W, Boeree MJ, van Soolingen D. Drug treatment of pulmonary nontuberculous mycobacterial disease in HIV-negative patients: the evidence. *Expert Rev Anti Infect Ther* 2013; **11**: 1065–77.
- Shen GH, Wu BD, Hu ST, Lin CF, Wu KM, Chen JH. High efficacy of clofazimine and its synergistic effect with amikacin against rapidly growing mycobacteria. *Int J Antimicrob Agents* 2010; **35**: 400–04.
- Brown-Elliott BA, Nash KA, Wallace RJ Jr. Antimicrobial susceptibility testing, drug resistance mechanisms, and therapy of infections with nontuberculous mycobacteria. *Clin Microbiol Rev* 2012; **25**: 545–82.
- Nash KA, Brown-Elliott BA, Wallace RJ Jr. A novel gene, erm(41), confers inducible macrolide resistance to clinical isolates of *Mycobacterium abscessus* but is absent from *Mycobacterium chelonae*. *Antimicrob Agents Chemother* 2009; **53**: 1367–76.
- Wallace RJ Jr, Meier A, Brown BA, et al. Genetic basis for clarithromycin resistance among isolates of *Mycobacterium chelonae* and *Mycobacterium abscessus*. *Antimicrob Agents Chemother* 1996; **40**: 1676–81.
- George IA, Santos CA, Olsen MA, Bailey TC. Epidemiology and outcomes of nontuberculous mycobacterial infections in solid organ transplant recipients at a midwestern center. *Transplantation* 2016; **100**: 1073–78.
- Goto T, Hamaguchi R, Maeshima A, Oyamada Y, Kato R. Pulmonary resection for *Mycobacterium chelonae* infection. *Ann Thorac Cardiovasc Surg* 2012; **18**: 128–31.
- Knoll BM, Kappagoda S, Gill RR, et al. Non-tuberculous mycobacterial infection among lung transplant recipients: a 15-year cohort study. *Transpl Infect Dis* 2012; **14**: 452–60.
- Ko Y, Kim W, Shin BS, et al. Nontuberculous mycobacterial lung disease caused by *Mycobacterium chelonae*: A Case Report. *Tuberc Respir Dis (Seoul)* 2013; **74**: 191–94.
- Peres E, Khaled Y, Krijanovski OI, et al. *Mycobacterium chelonae* necrotizing pneumonia after allogeneic hematopoietic stem cell transplant: report of clinical response to treatment with tigecycline. *Transpl Infect Dis* 2009; **11**: 57–63.
- Satyanarayana G, Heysell SK, Scully KW, Houpt ER. Mycobacterial infections in a large Virginia hospital, 2001-2009. *BMC Infect Dis* 2011; **11**: 113.
- Tsukamura M, Nakamura E, Kurita I, Nakamura T. Isolation of *Mycobacterium chelonae* subspecies *chelonae* (*Mycobacterium borstelense*) from pulmonary lesions of 9 patients. *Am Rev Respir Dis* 1973; **108**: 683–85.
- Al Jarad N, Demertzis P, Jones DJ, et al. Comparison of characteristics of patients and treatment outcome for pulmonary non-tuberculous mycobacterial infection and pulmonary tuberculosis. *Thorax* 1996; **51**: 137–39.
- Gupta N, Mittal A, Niyas VKM, et al. Nontuberculous mycobacteria: a report of eighteen cases from a tertiary care center in India. *Lung India* 2020; **37**: 495–500.
- Hadjiliadis D, Adlakha A, Prakash UB. Rapidly growing mycobacterial lung infection in association with esophageal disorders. *Mayo Clin Proc* 1999; **74**: 45–51.
- Huang CW, Chen JH, Hu ST, et al. Synergistic activities of tigecycline with clarithromycin or amikacin against rapidly growing mycobacteria in Taiwan. *Int J Antimicrob Agents* 2013; **41**: 218–23.
- McCallum AD, Watkin SW, Faccenda JF. Non-tuberculous mycobacterial infections in the Scottish borders: identification, management and treatment outcomes—a retrospective review. *J R Coll Physicians Edinb* 2011; **41**: 294–303.
- Ruas RR, Abreu I, Nuak J, et al. Nontuberculous mycobacteria in a tertiary hospital in Portugal: A clinical review. *Int J Mycobacteriol* 2017; **6**: 344–48.

- 32 Shah SK, McAnally KJ, Seoane L, et al. Analysis of pulmonary non-tuberculous mycobacterial infections after lung transplantation. *Transpl Infect Dis* 2016; **18**: 585–91.
- 33 Wallace RJ Jr, Dukart G, Brown-Elliott BA, Griffith DE, Scerpella EG, Marshall B. Clinical experience in 52 patients with tigecycline-containing regimens for salvage treatment of *Mycobacterium abscessus* and *Mycobacterium chelonae* infections. *J Antimicrob Chemother* 2014; **69**: 1945–53.
- 34 Yano Y, Kitada S, Mori M, et al. Pulmonary disease caused by rapidly growing mycobacteria: a retrospective study of 44 cases in Japan. *Respiration* 2013; **85**: 305–11.
- 35 Park S, Suh GY, Chung MP, et al. Clinical significance of *Mycobacterium fortuitum* isolated from respiratory specimens. *Respir Med* 2008; **102**: 437–42.
- 36 Nash KA, Zhang Y, Brown-Elliott BA, Wallace RJ Jr. Molecular basis of intrinsic macrolide resistance in clinical isolates of *Mycobacterium fortuitum*. *J Antimicrob Chemother* 2005; **55**: 170–77.
- 37 Wallace RJ Jr, Bedsole G, Sumter G, et al. Activities of ciprofloxacin and ofloxacin against rapidly growing mycobacteria with demonstration of acquired resistance following single-drug therapy. *Antimicrob Agents Chemother* 1990; **34**: 65–70.
- 38 Abe Y, Nakamura M, Suzuki K, et al. Massive hemoptysis due to *Mycobacterium fortuitum* infection controlled with bronchial artery embolization - a case report. *Clin Imaging* 1999; **23**: 361–63.
- 39 Al-Ghaffi H, Al-Hajj S. Nontuberculous mycobacteria in Saudi Arabia and Gulf Countries: a review. *Can Respir J* 2017; **2017**: 5035932.
- 40 Aronchick JM, Miller WT, Epstein DM, Gefter WB. Association of achalasia and pulmonary *Mycobacterium fortuitum* infection. *Radiology* 1986; **160**: 85–86.
- 41 Banerjee R, Hall R, Hughes GR. Pulmonary *Mycobacterium fortuitum* infection in association with achalasia of the oesophagus. Case report and review of the literature. *Br J Dis Chest* 1970; **64**: 112–18.
- 42 Bouchentouf R. *Mycobacterium fortuitum* infection associated with achalasia. *Tunis Med* 2018; **96**: 311–13.
- 43 Burns DN, Rohatgi PK, Rosenthal R, Seiler M, Gordin FM. Disseminated *Mycobacterium fortuitum* successfully treated with combination therapy including ciprofloxacin. *Am Rev Respir Dis* 1990; **142**: 468–70.
- 44 Chen CY, Sheng WH, Lai CC, et al. Mycobacterial infections in adult patients with hematological malignancy. *Eur J Clin Microbiol Infect Dis* 2012; **31**: 1059–66.
- 45 Corpe RF, Smith CE, Stergus I. Death due to *Mycobacterium fortuitum*. *JAMA* 1961; **177**: 262–63.
- 46 de Mello KG, Mello FC, Borga L, et al. Clinical and therapeutic features of pulmonary nontuberculous mycobacterial disease, Brazil, 1993–2011. *Emerg Infect Dis* 2013; **19**: 393–99.
- 47 Del Giudice G, Iadevaia C, Santoro G, Moscariele E, Smeraglia R, Marzo C. Nontuberculous mycobacterial lung disease in patients without HIV infection: a retrospective analysis over 3 years. *Clin Respir J* 2011; **5**: 203–10.
- 48 Diel R, Ringshausen F, Richter E, Welker L, Schmitz J, Nienhaus A. Microbiological and clinical outcomes of treating non-*Mycobacterium avium* complex nontuberculous mycobacterial pulmonary disease: a systematic review and meta-analysis. *Chest* 2017; **152**: 120–42.
- 49 Ellender CM, Law DB, Thomson RM, Eather GW. Safety of IV amikacin in the treatment of pulmonary non-tuberculous mycobacterial disease. *Respirology* 2016; **21**: 357–62.
- 50 Garcia-Ibarbia C, Espina B, Fernandez-Ayala M, Nan D. *Mycobacterium fortuitum* infection and lipoid pneumonia. *Respiratory Medicine Extra* 2007; **3**: 92–93.
- 51 Han XY, Dé I, Jacobson KL. Rapidly growing mycobacteria: clinical and microbiologic studies of 115 cases. *Am J Clin Pathol* 2007; **128**: 612–21.
- 52 Hasan A, Swamy TLN. Nocardia and *Mycobacterium fortuitum* infection in a case of lipoid pneumonia. *Respir Med CME* 2011; **4**: 75–78.
- 53 Howard RS 2nd, Woodring JH, Vandiviere HM, Dillon ML. *Mycobacterium fortuitum* pulmonary infection complicating achalasia. *South Med J* 1991; **84**: 1391–93.
- 54 Ichijima S, Tsukamura M. Ofloxacin and the treatment of pulmonary disease due to *Mycobacterium fortuitum*. *Chest* 1987; **92**: 1110–12.
- 55 Keating MR, Daly JS. Nontuberculous mycobacterial infections in solid organ transplantation. *Am J Transplant* 2013; **13** (suppl 4): 77–82.
- 56 Kupeli E, Bozkurt E, Azap O, Eyuboglu FO. *Mycobacterium fortuitum* infection presenting as community-acquired pneumonia in an immunocompetent host. *J Bronchology Interv Pulmonol* 2010; **17**: 356–58.
- 57 Kurokawa K, Harada N, Sasano H, et al. Pulmonary infection due to fluoroquinolone-resistant *Mycobacterium fortuitum*: a case report. *BMC Infect Dis* 2020; **20**: 866.
- 58 Lessing MP, Walker MM. Fatal pulmonary infection due to *Mycobacterium fortuitum*. *J Clin Pathol* 1993; **46**: 271–72.
- 59 Lopez-Luis BA, Sifuentes-Osorio J, Pérez-Gutiérrez MT, Chávez-Mazari B, Bobadilla-Del-Valle M, Ponce-de-León A. Nontuberculous mycobacterial infection in a tertiary care center in Mexico, 2001–2017. *Braz J Infect Dis* 2020; **24**: 213–20.
- 60 Martiniano SL, Wagner BD, Levin A, Nick JA, Sagel SD, Daley CL. Safety and effectiveness of clofazimine for primary and refractory nontuberculous mycobacterial infection. *Chest* 2017; **152**: 800–09.
- 61 Matsumoto T, Otsuka K, Tomii K. *Mycobacterium fortuitum* thoracic empyema: a case report and review of the literature. *J Infect Chemother* 2015; **21**: 747–50.
- 62 Napolitano C, Arunabh X, Mojaverian A, Shah R, Kaplan MR. *Mycobacterium fortuitum* pulmonary infection complicating achalasia. *Infect Dis Clin Pract* 2004; **12**: 178–80.
- 63 Okamori S, Asakura T, Nishimura T, et al. Natural history of *Mycobacterium fortuitum* pulmonary infection presenting with migratory infiltrates: a case report with microbiological analysis. *BMC Infect Dis* 2018; **18**: 1.
- 64 Panagiotou M, Papaioannou AI, Kostikas K, et al. The epidemiology of pulmonary nontuberculous mycobacteria: data from a general hospital in Athens, Greece, 2007–2013. *Pulm Med* 2014; **2014**: 894976.
- 65 Radzniwan MR, Tohid H, Ahmad S, Mohd AF, Md Anshar F. Isolation of *Mycobacterium fortuitum* in sputum specimens of a patient with chronic cough: Is it clinically significant? *Malays Fam Physician* 2014; **9**: 38–41.
- 66 Simons S, van Ingen J, Hsueh PR, et al. Nontuberculous mycobacteria in respiratory tract infections, eastern Asia. *Emerg Infect Dis* 2011; **17**: 343–49.
- 67 Vadakekalam J, Ward MJ. *Mycobacterium fortuitum* lung abscess treated with ciprofloxacin. *Thorax* 1991; **46**: 737–38.
- 68 Varghese G, Shepherd R, Watt P, Bruce JH. Fatal infection with *Mycobacterium fortuitum* associated with oesophageal achalasia. *Thorax* 1988; **43**: 151–52.
- 69 Wang CC, Lin MC, Liu JW, Wang YH. Nontuberculous mycobacterial lung disease in southern Taiwan. *Chang Gung Med J* 2009; **32**: 499–508.
- 70 Yu JA, Weyant MJ, Mitchell JD. Surgical treatment of atypical mycobacterial infections. *Thorac Surg Clin* 2012; **22**: 277–85.
- 71 Annobil SH, Jamjoom GA, Bobo R, Iyengar J. Fatal lipoid pneumonia in an infant complicated by *Mycobacterium fortuitum* infection. *Trop Geogr Med* 1992; **44**: 160–64.
- 72 Kim SY, Moon SM, Jhun BW, et al. Species distribution and macrolide susceptibility of *Mycobacterium fortuitum* complex clinical isolates. *Antimicrob Agents Chemother* 2019; **63**: e02331-18.
- 73 Rammaert B, Couderc LJ, Rivaud E, et al. *Mycobacterium genavense* as a cause of subacute pneumonia in patients with severe cellular immunodeficiency. *BMC Infect Dis* 2011; **11**: 311.
- 74 Mahmood M, Ajmal S, Abu Saleh OM, Bryson A, Marcelin JR, Wilson JW. *Mycobacterium genavense* infections in non-HIV immunocompromised hosts: a systematic review. *Infect Dis (Lond)* 2018; **50**: 329–39.
- 75 Hoefsloot W, van Ingen J, Peters EJ, et al. *Mycobacterium genavense* in the Netherlands: an opportunistic pathogen in HIV and non-HIV immunocompromised patients. An observational study in 14 cases. *Clin Microbiol Infect* 2013; **19**: 432–37.
- 76 Böttger EC, Teske A, Kirschner P, et al. Disseminated “*Mycobacterium genavense*” infection in patients with AIDS. *Lancet* 1992; **340**: 76–80.
- 77 Blanco Pérez JJ, Pérez González A, Morano Amado LE, et al. Clinical significance of environmental mycobacteria isolated from respiratory specimens of patients with and without silicosis. *Arch Bronconeumol* 2016; **52**: 145–50.

- 78 Bogdan C, Kern P, Richter E, et al. Systemic infection with *Mycobacterium genavense* following immunosuppressive therapy in a patient who was seronegative for human immunodeficiency virus. *Clin Infect Dis* 1997; **24**: 1245–47.
- 79 Doggett JS, Strasfeld L. Disseminated *Mycobacterium genavense* with pulmonary nodules in a kidney transplant recipient: case report and review of the literature. *Transpl Infect Dis* 2011; **13**: 38–43.
- 80 Charles P, Lortholary O, Dechartres A, et al. *Mycobacterium genavense* infections: a retrospective multicenter study in France, 1996–2007. *Medicine (Baltimore)* 2011; **90**: 223–30.
- 81 Böttger EC. *Mycobacterium genavense*: an emerging pathogen. *Eur J Clin Microbiol Infect Dis* 1994; **13**: 932–36.
- 82 Gaynor CD, Clark RA, Koontz FP, Emler S, Hirschel B, Schlesinger LS. Disseminated *Mycobacterium genavense* infection in two patients with AIDS. *Clin Infect Dis* 1994; **18**: 455–57.
- 83 Thomsen VO, Dragsted UB, Bauer J, Fuursted K, Lundgren J. Disseminated infection with *Mycobacterium genavense*: a challenge to physicians and mycobacteriologists. *J Clin Microbiol* 1999; **37**: 3901–05.
- 84 Albrecht H, Rüschi-Gerdes S, Stellbrink HJ, Greten H, Jäckle S. Disseminated *Mycobacterium genavense* infection as a cause of pseudo-Whipple's disease and sclerosing cholangitis. *Clin Infect Dis* 1997; **25**: 742–43.
- 85 Szturmowicz M, Siemion-Szcześniak I, Wyrostkiewicz D, et al. Factors predisposing to non-tuberculous mycobacterial lung disease in the patients with respiratory isolates of non-tuberculous mycobacteria. *Adv Respir Med* 2018; **86**: 261–67.
- 86 Hoefsloot W, van Ingen J, Andrejak C, et al. The geographic diversity of nontuberculous mycobacteria isolated from pulmonary samples: an NTM-NET collaborative study. *Eur Respir J* 2013; **42**: 1604–13.
- 87 Moore JE, Kruijshaar ME, Ormerod LP, Drobniewski F, Abubakar I. Increasing reports of non-tuberculous mycobacteria in England, Wales and Northern Ireland, 1995–2006. *BMC Public Health* 2010; **10**: 612.
- 88 Waak MB, LaPara TM, Hallé C, Hozalski RM. Nontuberculous mycobacteria in two drinking water distribution systems and the role of residual disinfection. *Environ Sci Technol* 2019; **53**: 8563–73.
- 89 Oriani AS, Marfil MJ, Zumárraga MJ, Baldini MD. Prevalence and species diversity of nontuberculous mycobacteria in drinking water supply system of Bahía Blanca City, Argentina. *Int J Mycobacteriol* 2019; **8**: 138–45.
- 90 Bauernfeind A. Comparative in-vitro activities of the new quinolone, Bay y 3118, and ciprofloxacin, sparfloracin, tosufloxacin, CI-960 and CI-990. *J Antimicrob Chemother* 1993; **31**: 505–22.
- 91 Goswami B, Narang P, Mishra PS, Narang R, Narang U, Mendiratta DK. Drug susceptibility of rapid and slow growing non-tuberculous mycobacteria isolated from symptomatics for pulmonary tuberculosis, Central India. *Indian J Med Microbiol* 2016; **34**: 442–47.
- 92 Brown-Elliott BA, Crist CJ, Mann LB, Wilson RW, Wallace RJ Jr. In vitro activity of linezolid against slowly growing nontuberculous mycobacteria. *Antimicrob Agents Chemother* 2003; **47**: 1736–38.
- 93 Craig CP, Kreitzer SM. Non-tuberculous mycobacterial infections: human infections due to *Mycobacterium gordonae*. *Infect Dis Rep* 1980; **6**: 79–88.
- 94 Chang HY, Tsai WC, Lee TF, Sheng WH. *Mycobacterium gordonae* infection in immunocompromised and immunocompetent hosts: a series of seven cases and literature review. *J Formos Med Assoc* 2021; **120**: 524–32.
- 95 de Gracia J, Vidal R, Martin N, Bravo C, Gonzalez T, Riba A. Pulmonary disease caused by *Mycobacterium gordonae*. *Tubercle* 1989; **70**: 135–37.
- 96 Douglas JG, Calder MA, Choo-Kang YF, Leitch AG. *Mycobacterium gordonae*: a new pathogen? *Thorax* 1986; **41**: 152–53.
- 97 Morimoto K, Kazumi Y, Shiraiishi Y, et al. Clinical and microbiological features of definite *Mycobacterium gordonae* pulmonary disease: the establishment of diagnostic criteria for low-virulence mycobacteria. *Trans R Soc Trop Med Hyg* 2015; **109**: 589–93.
- 98 Nanda Kumar U, Varkey B. Pulmonary infection caused by *Mycobacterium gordonae*. *Br J Dis Chest* 1980; **74**: 189–92.
- 99 Nathan V, Mehta JB, Dralle W. Rifabutin in the treatment of cavitary lung disease due to *Mycobacterium gordonae*. *South Med J* 1993; **86**: 839–41.
- 100 Umeda Y, Matsuno Y, Imaizumi M, Mori Y, Iwata H, Takiya H. Extralobar pulmonary sequestration infected with *Mycobacterium gordonae*. *J Thorac Cardiovasc Surg* 2009; **137**: e23–24.
- 101 Schroder KH, Juhlin I. *Mycobacterium malmoeense* sp. nov. *Int J Syst Bacteriol* 1977; **26**: 409.
- 102 Hoefsloot W, van Ingen J, de Lange WC, Dekhuijzen PN, Boeree MJ, van Soolingen D. Clinical relevance of *Mycobacterium malmoeense* isolation in the Netherlands. *Eur Respir J* 2009; **34**: 926–31.
- 103 Henry MT, Inamdar L, O'Riordain D, Schweiger M, Watson JP. Nontuberculous mycobacteria in non-HIV patients: epidemiology, treatment and response. *Eur Respir J* 2004; **23**: 741–46.
- 104 Research Committee of the British Thoracic Society. First randomised trial of treatments for pulmonary disease caused by *M avium intracellulare*, *M malmoeense*, and *M xenopi* in HIV negative patients: rifampicin, ethambutol and isoniazid versus rifampicin and ethambutol. *Thorax* 2001; **56**: 167–72.
- 105 British Thoracic Society. Pulmonary disease caused by *M. malmoeense* in HIV negative patients: 5-yr follow-up of patients receiving standardised treatment. *Eur Respir J* 2003; **21**: 478–82.
- 106 van Ingen J, van der Laan T, Dekhuijzen R, Boeree M, van Soolingen D. In vitro drug susceptibility of 2275 clinical non-tuberculous mycobacterium isolates of 49 species in the Netherlands. *Int J Antimicrob Agents* 2010; **35**: 169–73.
- 107 Engervall P, Björkholm M, Petrini B, Heurlin N, Henriques B, Källén G. Disseminated *Mycobacterium malmoeense* infection in a patient with chronic granulocytic leukaemia. *J Intern Med* 1993; **234**: 231–33.
- 108 Saito H, Tomioka H, Sato K, Tasaka H, Dekio S. *Mycobacterium malmoeense* isolated from soil. *Microbiol Immunol* 1994; **38**: 313–15.
- 109 Thorel MF, Falkinham JO 3rd, Moreau RG. Environmental mycobacteria from alpine and subalpine habitats. *FEMS Microbiol Ecol* 2004; **49**: 343–47.
- 110 Evans AJ, Crisp AJ, Colville A, Evans SA, Johnston ID. Pulmonary infections caused by *Mycobacterium malmoeense* and *Mycobacterium tuberculosis*: comparison of radiographic features. *AJR Am J Roentgenol* 1993; **161**: 733–37.
- 111 France AJ, McLeod DT, Calder MA, Seaton A. *Mycobacterium malmoeense* infections in Scotland: an increasing problem. *Thorax* 1987; **42**: 593–95.
- 112 Hoefsloot W, Boeree MJ, van Ingen J, et al. The rising incidence and clinical relevance of *Mycobacterium malmoeense*: a review of the literature. *Int J Tuberc Lung Dis* 2008; **12**: 987–93.
- 113 Banks J, Jenkins PA, Smith AP. Pulmonary infection with *Mycobacterium malmoeense*—a review of treatment and response. *Tubercle* 1985; **66**: 197–203.
- 114 Murray MP, Laurenson IF, Hill AT. Outcomes of a standardized triple-drug regimen for the treatment of nontuberculous mycobacterial pulmonary infection. *Clin Infect Dis* 2008; **47**: 222–24.
- 115 Jenkins PA, Campbell IA, Banks J, Gelder CM, Prescott RJ, Smith AP. Clarithromycin vs ciprofloxacin as adjuncts to rifampicin and ethambutol in treating opportunist mycobacterial lung diseases and an assessment of *Mycobacterium vaccae* immunotherapy. *Thorax* 2008; **63**: 627–34.
- 116 van Ingen J, Boeree MJ, Dekhuijzen PN, van Soolingen D. Clinical relevance of *Mycobacterium simiae* in pulmonary samples. *Eur Respir J* 2008; **31**: 106–09.
- 117 Rynkiewicz DL, Cage GD, Butler WR, Ampel NM. Clinical and microbiological assessment of *Mycobacterium simiae* isolates from a single laboratory in southern Arizona. *Clin Infect Dis* 1998; **26**: 625–30.
- 118 Valero G, Peters J, Jorgensen JH, Graybill JR. Clinical isolates of *Mycobacterium simiae* in San Antonio, Texas. An 11-yr review. *Am J Respir Crit Care Med* 1995; **152**: 1555–57.
- 119 Jabbour JF, Hamieh A, Sharara SL, Kanj SS. *Mycobacterium simiae*: harmless colonizer or deadly pathogen? *PLoS Pathog* 2020; **16**: e1008418.
- 120 Braun E, Sprecher H, Davidson S, Kassis I. Epidemiology and clinical significance of non-tuberculous mycobacteria isolated from pulmonary specimens. *Int J Tuberc Lung Dis* 2013; **17**: 96–99.
- 121 Shitrit D, Peled N, Bishara J, et al. Clinical and radiological features of *Mycobacterium kansasii* infection and *Mycobacterium simiae* infection. *Respir Med* 2008; **102**: 1598–603.

- 122 Hamieh A, Tayyar R, Tabaja H, et al. Emergence of *Mycobacterium simiae*: A retrospective study from a tertiary care center in Lebanon. *PLoS One* 2018; **13**: e0195390.
- 123 Nasiri MJ, Dabiri H, Darban-Sarokhalil D, Hashemi Shahraki A. Prevalence of non-tuberculosis mycobacterial infections among tuberculosis suspects in Iran: systematic review and meta-analysis. *PLoS One* 2015; **10**: e0129073.
- 124 Shenai S, Rodrigues C, Mehta A. Time to identify and define non-tuberculous mycobacteria in a tuberculosis-endemic region. *Int J Tuberc Lung Dis* 2010; **14**: 1001–08.
- 125 van Ingen J, Totten SE, Heifets LB, Boeree MJ, Daley CL. Drug susceptibility testing and pharmacokinetics question current treatment regimens in *Mycobacterium simiae* complex disease. *Int J Antimicrob Agents* 2012; **39**: 173–76.
- 126 Pang H, Jiang Y, Wan K. Drug susceptibility of 33 reference strains of slowly growing mycobacteria to 19 antimicrobial agents. *BioMed Res Int* 2017; **2017**: 1584658.
- 127 Cowman S, Burns K, Benson S, Wilson R, Loebinger MR. The antimicrobial susceptibility of non-tuberculous mycobacteria. *J Infect* 2016; **72**: 324–31.
- 128 Baghaei P, Tabarsi P, Farnia P, et al. Pulmonary disease caused by *Mycobacterium simiae* in Iran's national referral center for tuberculosis. *J Infect Dev Ctries* 2012; **6**: 23–28.
- 129 Coolen-Allou N, Touron T, Belmonte O, et al. Clinical, radiological, and microbiological characteristics of *Mycobacterium simiae* infection in 19 patients. *Antimicrob Agents Chemother* 2018; **62**: e00395-18.
- 130 Hashemi-Shahraki A, Darban-Sarokhalil D, Heidarieh P, et al. *Mycobacterium simiae*: a possible emerging pathogen in Iran. *Jpn J Infect Dis* 2013; **66**: 475–79.
- 131 Jeong SH, Kim SY, Lee H, et al. Nontuberculous mycobacterial lung disease caused by *Mycobacterium simiae*: the first reported case in South Korea. *Tuberc Respir Dis (Seoul)* 2015; **78**: 432–35.
- 132 Aharon A, Langevitz P, Maran R, Blank-Porat D, Shtrasburg S, Livneh A. Reactive amyloidosis in a patient with *Mycobacterium simiae* pulmonary infection. *Respir Med* 1998; **92**: 123–24.
- 133 Rose HD, Dorff GJ, Lauwasser M, Sheth NK. Pulmonary and disseminated *Mycobacterium simiae* infection in humans. *Am Rev Respir Dis* 1982; **126**: 1110–13.
- 134 Reddy LM, Bhaskar H, Prashanth N, Kumar Narahari NK, Paramjyothi GK. All Mycobacteria are not tubercular: A case report. *J Clin Diagn Res* 2018; **12**: OD01–03.
- 135 McSweeney FG, O'Brien ME, Sheehan S, Plant B, Corcoran G. Pulmonary *Mycobacterium szulgai* infection. *Ir Med J* 2012; **105**: 275–77.
- 136 van Ingen J, Boeree MJ, de Lange WC, de Haas PE, Dekhuijzen PN, van Soolingen D. Clinical relevance of *Mycobacterium szulgai* in the Netherlands. *Clin Infect Dis* 2008; **46**: 1200–05.
- 137 Maloney JM, Gregg CR, Stephens DS, Manian FA, Rimland D. Infections caused by *Mycobacterium szulgai* in humans. *Rev Infect Dis* 1987; **9**: 1120–26.
- 138 Schaefer WB, Wolinsky E, Jenkins PA, Marks J. *Mycobacterium szulgai*-a new pathogen. Serologic identification and report of five new cases. *Am Rev Respir Dis* 1973; **108**: 1320–26.
- 139 Davidson PT. *Mycobacterium szulgai*; a new pathogen causing infection of the lung. *Chest* 1976; **69**: 799–801.
- 140 Medinger AE, Spagnolo SV. *Mycobacterium szulgai* pulmonary infection: the importance of knowing. *South Med J* 1981; **74**: 85–86.
- 141 Collazos J, Díaz F, Rodríguez J, Ayarza R. Persistent lung infection due to *Mycobacterium szulgai*. *Tuberc Lung Dis* 1993; **74**: 412–13.
- 142 Benator DA, Kan V, Gordin FM. *Mycobacterium szulgai* infection of the lung: case report and review of an unusual pathogen. *Am J Med Sci* 1997; **313**: 346–51.
- 143 Tortoli E, Besozzi G, Lacchini C, Penati V, Simonetti MT, Emler S. Pulmonary infection due to *Mycobacterium szulgai*, case report and review of the literature. *Eur Respir J* 1998; **11**: 975–77.
- 144 Tsuyuguchi K, Amitani R, Matsumoto H, et al. A resected case of *Mycobacterium szulgai* pulmonary disease. *Int J Tuberc Lung Dis* 1998; **2**: 258–60.
- 145 Nakayama S, Fujii T, Kadota J, et al. Pulmonary mycobacteriosis caused by rifampicin-resistant *Mycobacterium szulgai*. *Intern Med* 2000; **39**: 309–12.
- 146 Tsuneyuka Y, Sato H, Hiranuma C. Surgical outcome of mycobacterium other than *Mycobacterium tuberculosis* pulmonary disease. *Thorac Cardiovasc Surg* 2000; **48**: 290–93.
- 147 Sánchez-Alarcos JM, De Miguel-Díez J, Bonilla I, Sicilia JJ, Alvarez-Sala JL. Pulmonary infection due to *Mycobacterium szulgai*. *Respiration* 2003; **70**: 533–36.
- 148 Hotta M, Minami Y, Itoda I, Yoshimori K, Takano K. A young female patient with anorexia nervosa complicated by *Mycobacterium szulgai* pulmonary infection. *Int J Eat Disord* 2004; **35**: 115–19.
- 149 Gutierrez M, Feola M, Lenge L, Rey R, Hoffman M. First pulmonary case reported in Argentina of infection with *mycobacterium szulgai*, a rare pathogen. *J Clin Microbiol* 2007; **45**: 3121–24.
- 150 Peloquin CA, Berning SE, Nitta AT, et al. Aminoglycoside toxicity: daily versus thrice-weekly dosing for treatment of mycobacterial diseases. *Clin Infect Dis* 2004; **38**: 1538–44.

© 2022 Elsevier Ltd. All rights reserved.